



Inactivation of vhsv by infiltration and salt under experimental conditions

Skall, H. F.; Jørgensen, C.; Olesen, Niels Jørgen

Published in:
17th International Conference on Diseases of Fish and Shellfish

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Skall, H. F., Jørgensen, C., & Olesen, N. J. (2015). Inactivation of vhsv by infiltration and salt under experimental conditions. In *17th International Conference on Diseases of Fish and Shellfish* (pp. 149-149)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

INACTIVATION OF VHSV BY INFILTRATION AND SALT UNDER EXPERIMENTAL CONDITIONS

H.F. SKALL^{1,3*}, C. JØRGENSEN², N.J. OLESEN³

¹*Aarhus University, Aarhus, Denmark*

²*DHI Hørsholm, Denmark*

³*National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark*

At the moment the only legal method in Denmark to sanitize wastewater from fish cutting plants is by infiltration. To evaluate the inactivation effect of infiltration on VHSV an experimental examination was initiated. A column packed with gravel as top- and bottom layer (total of 22 cm) and a mid layer consisting of dug sand (76 cm) was used for the trial. Over a period of 18 h 3.9×10^{10} TCID₅₀ VHSV was supplied to the column, where after normal tap water was supplied for the rest of the trial period, in total 7 days. During the 7 days samples for virological examination were taken. The sampling was most intensive in the period where the risk of VHSV breaking through the column was highest. The sensitivity of the virological examination was 13.9 TCID₅₀/ml and no virus was isolated. A reduction of VHSV > 4 log in the outlet water was seen. This experiment suggests that infiltration can be a valuable method to sanitize VHSV infected water. Changes in temperature, pH, earth types in the area used for infiltration etc. may change the virus reduction, though.

As some of the fish cutting plants are also smoking rainbow trout fillets, the question arose whether a brine solution will inactivate VHSV. In order to answer this question a small trial was set up. VHSV and NaCl was added to cell culture medium with 10% foetal bovine serum, in order to mimic a "dirty" environment, to obtain from 1.9% to 20.9% NaCl and kept in the dark at 4°C. Samples were titrated after 5 min, 1 h and 20 h. No reduction in titre was observed in any of the samples.